AD			

Award Number: DAMD17-01-1-0817

TITLE: Bone Growth, Mechanical Stimulus and IGF-I

PRINCIPAL INVESTIGATOR: Vicente Gilsanz, M.D.

CONTRACTING ORGANIZATION: Children's Hospital Los Angeles

Los Angeles, CA 90027

REPORT DATE: October 2007

TYPE OF REPORT: Addendum to Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

data needed, and completing a this burden to Department of D	and reviewing this collection of i refense, Washington Headquar	nformation. Send comments regaters Services, Directorate for Info	arding this burden estimate or an mation Operations and Reports	y other aspect of this coll (0704-0188), 1215 Jeffer	lection of information, including suggestions for reducing roon Davis Highway, Suite 1204, Arlington, VA 22202-
		y other provision of law, no persor IR FORM TO THE ABOVE ADDF		or failing to comply with	a collection of information if it does not display a currently
1. REPORT DATE (DE	D-MM-YYYY)	2. REPORT TYPE			ATES COVERED (From - To)
01-10-2007 4. TITLE AND SUBTIT		Addendum to Final			SEP 2006 - 10 SEP 2007
4. IIILE AND SUBIII	LE			5a. C	CONTRACT NUMBER
Pana Crowth Mac	haniaal Stimulus a	ndICEI		5h (GRANT NUMBER
bone Growth, wec	chanical Stimulus a	na iGr-i			MD17-01-1-0817
					PROGRAM ELEMENT NUMBER
6. AUTHOR(S)				5d. F	PROJECT NUMBER
Vicente Gilsanz, M	1.D.				
				5e. 1	TASK NUMBER
E-Mail: vgilsanz@	chla.usc.edu			5f. V	VORK UNIT NUMBER
7. PERFORMING ORG	SANIZATION NAME(S)	AND ADDRESS(ES)			ERFORMING ORGANIZATION REPORT UMBER
Children's Hospita	l I os Angeles			l N	UWIDER
Los Angeles, CA	ū				
LUS Aligeles, CA	30021				
9. SPONSORING / MC	NITORING AGENCY N	IAME(S) AND ADDRESS	S(ES)	10. 5	SPONSOR/MONITOR'S ACRONYM(S)
U.S. Army Medica			2(=0)	1010	
Fort Detrick, Maryl					
, ,				11. 9	SPONSOR/MONITOR'S REPORT
				1	NUMBER(S)
12. DISTRIBUTION / A	VAILABILITY STATE	MENT			
Approved for Publ	ic Release; Distribι	ıtion Unlimited			
13. SUPPLEMENTAR	YNOTES				
14. ABSTRACT	atudu waa ta aaaaa	the netential for brief n	ariada af law magnitus	da high fraguana	www.machaniaal atimulation aignala in the
					by mechanical stimulation signals in the signals, several orders of magnitude
					skeleton of young adult females with
					evelopment and decrease the risk for
stress fractures in military recruits. Moreover, should these musculoskeletal changes persist through adulthood, this intervention may prove					
a deterrent to osteoporosis in the elderly.					
15. SUBJECT TERMS					
Mechanical Intervention, Fractures, IGF-I, Teenagers, Low Bone Mass					
	,	.			
16. SECURITY CLASSIFICATION OF:			17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON
			OF ABSTRACT	OF PAGES	USAMRMC
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area
U U U		UU	8	code)	

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

Table of Contents

Introduction	5
Body	5
Key Research Accomplishments	6
Reportable Outcomes	6
Conclusions	7
Personnel	7
Bibliography	8
Appendices	

INTRODUCTION

The incidence of osteoporosis, a disease that manifests in the elderly, may be reduced by increasing peak bone mass in the young. Indeed, susceptibility for low bone mass is present early in life, the amount of bone gained during adolescence is a main contributor to peak bone mass in the young adult, and peak bone mass in the young adult is a likely determinant of osteoporosis in the elderly. While research continues to identify means of reversing osteoporosis in the elderly, these data from children, adolescents and young adults indicate that enhancing bone health early in life represents a viable means of deterring osteoporosis decades before it arises. However, the benefits of early pharmacological interventions to prevent a disease that will not manifest for decades must be weighed against the possible complications of extended treatment. Preliminary data indicate that extremely low-level mechanical signals are anabolic to bone tissue, and their ability to enhance bone and muscle mass in young women was investigated in this study.

BODY

This study was designed to establish if brief, daily exposure to extremely low-level mechanical stimuli is anabolic to musculoskeletal development in young males and females, 15-20 years of age, with low bone density, who had previously sustained a fracture.

The effects of two twelve-month interventions on musculoskeletal development in young men and women were longitudinally studied and the results compared to matched groups of subjects undergoing no intervention. The mechanical intervention consisted of brief exposure to low level (0.3g; 1g = earth gravitational field) high frequency (30-Hz) mechanical loading for 10 minutes every day. The resistance exercise intervention consisted of 30 minutes of weight-bearing and trunk stabilization exercises three times per week.

A twelve-month trial was conducted in 48 young women (15-20y) with low bone density and a history of at least one skeletal fracture. Half of the subjects underwent brief (10 minute requested), daily, low-level whole body vibration (30 Hz, 0.3g); the remaining women served as controls. Quantitative computed tomography (CT) performed at baseline and at the end of study was used to establish changes in muscle and bone mass in the weight-bearing skeleton.

Using an Intention to Treat (ITT) analysis, cancellous bone in the lumbar vertebrae and cortical bone in the femoral midshaft of the experimental group increased by 2.1% (p=0.025) and 3.4% (p<0.001), respectively, as compared to 0.1% (p=0.74) and 1.1% (p=0.14), in controls. Increases in cancellous and cortical bone were 2.0% (p=0.06) and 2.3% (p=0.04) greater, respectively, in the experimental group when compared with controls. Cross-sectional area of paraspinous musculature was 4.9% greater (p=0.002) in the experimental group versus controls. When a per protocol analysis was considered, gains in both muscle and bone were strongly correlated to a threshold in compliance, where the benefit of the mechanical intervention as compared to controls was realized once subjects used the device for at least two minutes per day (n=18), as reflected by a 3.9% increase in cancellous bone of the spine (p=0.007), 2.9% increase in cortical bone of the femur (p=0.009), and 7.2% increase in musculature of the spine (p=0.001), as compared to controls and low-compliers (n=30).

Additionally, 24 males were initially enrolled in each of the vibration intervention and control groups. During the course of the intervention, one participant in the vibration intervention group moved out of state and one was incarcerated, the remaining 22 completed the intervention. There was no obvious benefit of the intervention in this group of subjects. These measures included paraspinous musculature, spine cancellous bone mineral density, spine cross-sectional area, visceral fat, subcutaneous fat, total fat, vertebral height and vertebral volume in the axial skeleton, and quadriceps femoris area, femoral cross-sectional area, femoral cortical bone area, femoral bone mineral density and femoral fat in the appendicular skeleton. Unfortunately, males were less compliant than females, possibly accounting for the negative results of this study

Serum levels of IGF-I were examined prior to and following the mechanical intervention in both study subjects and controls. At the mid-shaft of the femurs, IGF-I did not correlate with the material density of cortical bone (r = -0.08), but did correlate significantly with cortical bone area (r = 0.50; P < 0.0001) and with the cross-sectional area (r = 0.49; P < 0.0001) of the bone. When using multiple regression analyses, IGF-I was associated with both the cross-sectional area (P = 0.03) and cortical bone area (P = 0.04), even after accounting for age, gender, weight and the length of the femur. Thus, in the appendicular skeleton of male and female teenagers and young adults in this study, IGF-I had no influence on the material density of the bone, but was found to be a major determinant of the cross-sectional properties of the bone.

KEY RESEARCH ACCOMPLISHMENT

Establishing that short bouts of a low-level mechanical signal increases bone and muscle mass in young adult females.

REPORTABLE OUTCOMES

Abstracts and Presentations

Liu, X., C. Rosen, S. Mora, D. Dong, P. Pitukcheewanont, and V. Gilsanz, Low DXA and CT bone measures in young adults with a simple sequence repeat in IGF-I gene. J Bone Min Res, 2004. 19: p. S248.

Shepherd, J.A., B. Fan, M. Sherman, et al., Pediatric DXA precision varies with age. J Bone Min Res, 2004. 19: p. S234.

Wren, T.A.L., V. Gilsanz, P. Pitukcheewanont, et al., The Bone Mineral Density in Childhood Study (BMDCS): Substudy Comparing DXA and CT Vertebral Bone Measurements in Healthy Children and Adolescents. J Bone Min Res, 2004. 19: p. S67.

10/2004 "Low DXA and CT Bone Measures in Young Adults with a Simple Sequence Repeat in IGF-I Gene" 26th Annual Meeting of the American Society for Bone and Mineral Research Seattle, WA

03/2005	"Comparison of CT and DXA Measurements in Healthy Children" Bone Mineral Density in Childhood Study (BMDCS) Meeting National Institute of Child Health and Human Development Bethesda, MD
09/2005	"Mechanical Intervention Enhances Bone and Muscle in Young Women with Low Bone Density" American Society of Bone and Mineral Research 27 th Annual Meeting
06/2006	"Mechanical Intervention Enhances Bone and Muscle in Young Women with Low Bone Density" CHLA Saban Research Institute 11 th Annual Poster Session
06/2006	"Fat Mass is Not Beneficial to Bone" CHLA Saban Research Institute 11 th Annual Poster Session
06/2006	"Assessment of Vertebral Peak Bone Mass by CT and DXA" CHLA Saban Research Institute 11 th Annual Poster Session
09/2006	"Good, Good, Good Vibrations: Evidence for the Therapeutic Potential of Low- Magnitude, High Frequency Mechanical Signals" American Society of Bone and Mineral Research 28 th Annual Meeting

CONCLUSION

Short bouts of extremely low-level mechanical signals, several orders of magnitude below that associated with vigorous exercise, increased bone and muscle mass in the weight bearing skeleton of young adult females with low bone density. Should these musculoskeletal enhancements be preserved through adulthood, this intervention may prove a deterrent to osteoporosis in the elderly.

PERSONNEL

The salaries of the following personnel were supported through this research work:

Elizabeth Anton Monique Sanchez
Noel Arugay Valerie Thompson
Jaime Gonzalez Lisa Villanueva
Agnieska Janicka Cara Wah

Montre Koh Vicente Gilsanz, M.D.

Mercedes Landaverde Pisit Pitukcheewanont, M.D.

Sandra Maravilla

BIBLIOGRAPHY

Liu, X., C. Rosen, S. Mora, D. Dong, P. Pitukcheewanont, and V. Gilsanz, Low DXA and CT bone measures in young adults with a simple sequence repeat in IGF-I gene. J Bone Min Res, 2004. 19: p. S248.

Shepherd, J.A., B. Fan, M. Sherman, et al., Pediatric DXA precision varies with age. J Bone Min Res, 2004. 19: p. S234.

Wren, T.A.L., V. Gilsanz, P. Pitukcheewanont, et al., The Bone Mineral Density in Childhood Study (BMDCS): Substudy Comparing DXA and CT Vertebral Bone Measurements in Healthy Children and Adolescents. J Bone Min Res, 2004. 19: p. S67.

Wren, T.A.L., X. Liu, P. Pitukcheewanont, and V. Gilsanz, Bone densitometry in pediatric populations: discrepancies in the diagnosis of osteoporosis by DXA and CT. J Peds, accepted for publication on 1/13/2005, 2004.

Sanchez, M. and V. Gilsanz, Pediatric DXA bone measurements, P.E. Review, Editor. 2005. p. 19-23.

Wren, T.A., X. Liu, P. Pitukcheewanont, and V. Gilsanz, Bone Acquisition in Healthy Children and Adolescents: Comparisons of DXA and CT Measures. J Clin Endocrinol Metab, 2005. 90: p. 1925-1928.

Gilsanz, V., T.A. Wren, M. Sanchez, F. Dorey, S. Judex, and C. Rubin, Low-level, high-frequency mechanical signals enhance musculoskeletal development of young women with low BMD. J Bone Miner Res, 2006. 21(9): p. 1464-74.

Wren, T.A. and V. Gilsanz, Assessing bone mass in children and adolescents. Curr Osteoporos Rep, 2006. 4(4): p. 153-8.

Gilsanz, V. and T. Wren, Assessment of bone acquisition in childhood and adolescence. Pediatrics, 2007. 119 Suppl 2: p. S145-9.

Kalkwarf, H.J., B.S. Zemel, V. Gilsanz, et al., The bone mineral density in childhood study: bone mineral content and density according to age, sex, and race. J Clin Endocrinol Metab, 2007. 92(6): p. 2087-99.

Wren, T.A., P.S. Kim, A. Janicka, M. Sanchez, and V. Gilsanz, Timing of peak bone mass: discrepancies between CT and DXA. J Clin Endocrinol Metab, 2007. 92(3): p. 938-41.

Wren, T.A., X. Liu, P. Pitukcheewanont, and V. Gilsanz, Bone acquisition in healthy children and adolescents: comparisons of dual-energy x-ray absorptiometry and computed tomography measures. J Clin Endocrinol Metab, 2005. 90(4): p. 1925-8.

Wren, T.A., X. Liu, P. Pitukcheewanont, and V. Gilsanz, Bone densitometry in pediatric populations: discrepancies in the diagnosis of osteoporosis by DXA and CT. J Pediatr, 2005. 146(6): p. 776-9.

Sanchez, M.M. and V. Gilsanz, Pediatric DXA bone measurements. Pediatr Endocrinol Rev, 2005. 2 Suppl 3: p. 337-41.

Klein, G.L., L.A. Fitzpatrick, C.B. Langman, et al., The state of pediatric bone: summary of the ASBMR pediatric bone initiative. J Bone Miner Res, 2005. 20(12): p. 2075-81.